

Dr. Han Wrist Ganglion Cyst Excision

GLOVES: 6.5 Micro Indicator/6.5 Micro

Anesthesia: Choice

PREP: Hibiclens

POSITION: SUPINE, Patient on Stretcher, Rolling Hand Table, Tourniquet, Extremity Drape, Sitting Stools me and my PA

MEDICATIONS: 0.25% BUPIVACAINE or 0.5% Plain BUPIVACAINE at the end of the case.
If Patient and Anesthesia Want to do MAC, Then Please have 1% Lidocaine with Epinephrine Available in The Room

INSTRUMENTS: Hand Kit (Curved Littler Scissors or Curved Tenotomy Scissors, Ragnel x2 and Sen Retractor x 2, Ragnel, Small Weitlaner and Heiss Retractor)

EQUIPMENT: Tourniquet

SUPPLIES: HALF SHEET
Extremity Drape
Ray-Tech
LIGHT HANDLE COVER
15 BLADE
10CC SYRINGE (1)
25 GAUGE HYPO (1)
BIPOLAR CORD
TOURNIQUET 18"
4" ESMARK (OPEN, STERILE)

SUTURES: 4-0 NYLON P-C 3 or 3.0 Monocryl (Ask), Steri Strips

DRESSINGS: Xeroform, WET 4X4, DRY 4X4, 4" cast padding, 4" ACE Bandage, 4" Plaster x 10 Sheet

NOTES: BIPOLAR 15, NO BOVIE,

LAST UPDATED: 1/29/2019