Dr. Han Hand Open Reduction & Internal Fixation and Percutaneous Pinning

GLOVES: 6.5 Micro Indicator/6.5 Micro

Anesthesia: by Choice

PREP: Hibclens

POSITION: SUPINE, Patient on Stretcher, Rolling Hand Table, Tourniquet with "10 x 10" around Tourniquet, Finger Tourniquet Available, Extremity Drape, Sitting Stools

MEDICATIONS: 0.9% NACL IRRIGATION 500 ML
1% LIDOCAINE if MAC and Local (Pre-Inject)
Otherwise 0.25 Marcaine or 0.5% Marcaine whichever is available at end of case.

INSTRUMENTS: Lead Hand or Tupper Universal Hand Holder, Hand Tray,

EQUIPMENT: Tourniquet, Mini-C arm (drape), Small Battery Driver, Pin Cutter, double ended k-wires. Stryker Hand Plating System **(Do not open)**

SUPPLIES:

HALF SHEET
Extremity Drape
Ray-Tech
LIGHT HANDLE COVER
15 BLADE (2)
10CC SYRINGE (1)
25 GAUGE HYPO (1)
BIPOLAR CORD
TOURNIQUET 18"
4" ESMARK (OPEN, STERILE)

SUTURES: Ask. Do Not Open. 4-0 Nylon, 3-0 Fiber Wire or 3.0 Fiber-Loop for Tendon if available, if not 3.0 Ethibond, 5.0 Chromatic, 4.0 Chromatic

DRESSINGS: ADAPTIC or Xeroform, WET 4X4, DRY 4X4, 4" WEBRIL or 4" Cast Padding , 3" or 4" PLASTER, 3"/4" ACE WRAP, SLING (IF BLOCKED)

NOTES: BIPOLAR 15, NO BOVIE, Micro-Instrument and GEM Clip Appliers and Microscope in case for vessel and nerve repairs (Do not open)

LAST UPDATED: 8/27/2018