

Dr. Han Skin Graft Split Thickness

GLOVES: 6.5 Micro Indicator/6.5 Micro

Anesthesia: General or By Choice

PREP: Hibclens. Chloprep for Donor Site

POSITION: SUPINE, Patient on Stretcher if possible, Rolling Hand Table, Tourniquet with "10 x 10" around Tourniquet, Extremity Drape, Sitting Stools

MEDICATIONS: 0.9% NAACL IRRIGATION
30 ML 1% LIDOCAINE with Epinephrine add to 70 ml of Injectable Saline (A solution to tumescent the skin graft donor site)
Mineral Oil

INSTRUMENTS: Lead Hand or Tupper Hand Retractor Do not Open, Plastic Surgery Tray

EQUIPMENT: Zimmer Dermatome, Dermatome Blade and 3 inch Guard, Zimmer Skin Mesher and 1 x 1.5 Skin Graft Carrier

SUPPLIES:

HALF SHEET
Skin Stapler
Extremity Drape
Ray-Tech
LIGHT HANDLE COVER
15 BLADE (2)
60 CC SYRINGE (1)
20 or 18 gauge Spinal Needle
TOURNIQUET 18"
Bovie
4" ESMARK (OPEN, STERILE)
4" WEBRIL (UNDER TOURNIQUET) if needed

SUTURES: 4.0-Chromatic SH
Stapler
3.0 Silk Pop Off Sutures -Do Not Open.

DRESSINGS: **Skin Graft:** 1. Adaptic and Wound Vac if requested or 2. Xeforum, Cotton Bolster (Ask), Mastisol, 4x4 Gauze, 4" Kerlex, and 4" ACE Wrap x 2
Donor Site: Xeroform, 4x4 Gauze, ABD Pads, 4" Kerlex, and 4" ACE Wrap x 2

NOTES: BOVIE and Bipolar

LAST UPDATED: 8/27/2018

